

NEWS RELEASE

1111 North Fairfax Street
Alexandria, VA 22314-1488
703 684 2782
703 684 7343 fax
www.apta.org

Physical Therapists Play Integral Role in Prevention, Risk Reduction, and Treatment of Painful and Often Irreversible Side Effect of Cancer Treatment

IN OBSERVANCE OF BREAST CANCER AWARENESS, PHYSICAL THERAPISTS PROMOTE EARLY DIAGNOSIS FOR SUCCESSFUL TREATMENT OF LYMPHEDEMA

FOR IMMEDIATE RELEASE

Contact: Erin Wendel
erinwendel@apta.org
703-706-3397

ALEXANDRIA, VA – As breast cancer awareness month is observed during October, the American Physical Therapy Association (APTA) is hoping to shine a spotlight on [lymphedema](#), a chronic, debilitating and often irreversible side effect of cancer treatment. According to APTA, breast cancer patients who seek the services of a physical therapist can reduce their risk of lymphedema, as it can be prevented or more effectively managed if caught in its earliest stages and treated by a physical therapist. APTA is launching this effort as it kicks off [National Physical Therapy Month](#) in October to educate the public about the important role physical therapists and physical therapist assistants play in health care.

Breast cancer-related lymphedema, which can cause significant swelling of the upper and lower extremities due to the build-up of excess lymph fluid, is mostly caused by damage to the body's lymphatic system during treatment for cancer and can include limited movement, joint pain, and difficulty performing activities. Physical therapists, who are experts in restoring motion and mobility in people's lives, play an important role not only in treating the effects of lymphedema but also in reducing the risk of occurrence.

A study published in the journal *Cancer* (April 25, 2008) showed that pre-operative assessments of patients with breast cancer by physical therapists allow for earlier diagnosis and treatment of lymphedema. This includes programs that use infra-red technology to measure limb volume and regular post-operative follow-up to detect lymphedema.

- more -

The study further demonstrated that when lymphedema was detected early, the patient's condition was managed with a more conservative approach that included a light-grade compression sleeve and hand gauntlet that effectively decreased early swelling and possibly prevented the progression of lymphedema. Fitted by a physical therapist, these were prescribed for daily wear.

"Lymphedema is normally treated with more aggressive and often costly and time-consuming techniques, such as complete decongestive therapy," explained APTA Board member and spokesperson, and the study's lead author, Nicole Stout, PT, MPT, CLT-LANA. Complete decongestive therapy requires the patient to attend daily therapy sessions for weeks and wear bulky compression bandages.

"This study clearly demonstrates the fact that lymphedema can be managed with a more conservative treatment option when diagnosed in its earliest stages, which is good news for patients with breast cancer," Stout said. "Based on these findings, we encourage patients with breast cancer to seek referral for physical therapy services before, during and after breast cancer treatment because arm problems can occur at any time during the course of care and the best outcomes are reached when they can be detected early, or even prevented.

"We hope physicians, surgeons, oncologists, and other physical therapists will make early intervention and conservative treatment of lymphedema the standard of care in breast cancer treatment," Stout concluded.

APTA Board member Babette Sanders, PT, MS, knows firsthand the benefits of using physical therapy for prevention and maintenance of keeping lymphedema at bay. "Swelling occurred immediately in my upper arm and trunk after my mastectomy," she said. "By quickly incorporating physical therapy as part of my overall management of the condition, I was able to get rid of the swelling and have since kept it away. I am able to remain lymphedema free with monthly preventative treatment. It makes a tremendous difference in the quality of my daily living and my ability to move and function."

Stout recommends patients with breast cancer be aware of their risk factors and indicators of the onset of lymphedema:

Risk Factors

- Being overweight or gaining weight during or after cancer treatment
- Having lymph nodes removed
- Having radiation therapy
- Having an infection or blood clot in the limb affected by cancer treatment

Warning Signs

- Aching or heaviness in the limb that was affected by cancer treatment
- A feeling of fullness or tightness in the limb affected by cancer treatment
- Visible swelling that affects the limb affected by cancer treatment

Some risk factors, such as the necessary treatment for cancer, cannot be avoided. However, there are many things you *can* do to help reduce the risk of lymphedema, such as:

- Maintain a healthy body weight
- Follow proper skin care and hygiene practices
- Avoid infections from cuts, burns, or skin wounds
- Exercise sensibly and avoid unnecessary strain to the limb

A physical therapist will create an exercise program tailored to your individual needs and monitor its effect on your limb.

#

Physical therapists are highly-educated, licensed health care professionals who can help patients reduce pain and improve or restore mobility — in many cases without expensive surgery or the side effects of prescription medications. APTA represents more than 74,000 physical therapists, physical therapist assistants, and students of physical therapy nationwide. Its purpose is to improve the health and quality of life of individuals through the advancement of physical therapist practice, education, and research. In most states, patients can make an appointment directly with a physical therapist, without a physician referral. Learn more about conditions physical therapists can treat and find a physical therapist in your area at www.moveforwardpt.com. Join APTA on [Facebook](#) and [Twitter](#).